

OBSTRUCTING SLICENESS IN A FAMILY OF MONTESINOS KNOTS

LUKE WILLIAMS

ABSTRACT. Using Gauge theoretical techniques employed by Lisca for 2-bridge knots and by Greene-Jabuka for 3-stranded pretzel knots, we show that no member of the family of Montesinos knots $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$, with certain restrictions on m_i , n_i , and q , can be (smoothly) slice. Our techniques use Donaldson's diagonalization theorem and the fact that the 2-fold covers of Montesinos knots bound plumbing 4-manifolds, many of which are negative definite. Some of our examples include knots with signature 0 and square determinant.

1. INTRODUCTION

In his recent breakthrough articles [Lis1, Lis2], Lisca applies gauge theory, based on work of Donaldson [Don], to obstruct smooth sliceness for 2-bridge knots. Lisca's approach uses the observation that the 2-fold branched cover of every 2-bridge knot bounds a negative definite plumbing 4-manifold. On the other hand, if a 2-bridge knot is slice, its 2-fold cover also bounds a rational homology 4-ball. Gluing these two 4-manifolds along their common boundary yields a smooth, closed, negative definite 4-manifold X . As such, according to Donaldson, its intersection form has to be standard. As Lisca goes on to show, this obstruction suffices to pin down all slice 2-bridge knots and moreover, suffices to determine the smooth concordance orders of all 2-bridge knots.

In [GJ], Greene and Jabuka use this approach, supplemented by Heegaard Floer homology techniques, to determine all slice knots among the 3-stranded pretzel knots $P(p, q, r)$ with p, q, r odd. Both Lisca's article, in the case of 2-bridge knots, and Greene-Jabuka's article, in the case of 3-stranded pretzel knots, resolve in the affirmative the slice-ribbon conjecture [Kir].

Building on Lisca's work, we employ the same gauge theoretic techniques to address the question of smooth sliceness for a five parameter family of Montesinos knots with three rational tangles. Specifically, the family we consider is (see Section 3 for a precise definition)

$$M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$$

with $m_i, n_i, -q \in \mathbb{N}$. Our main theorem is

Theorem 1.1. *Let $m_1, n_1, m_2, n_2, -q \in \mathbb{N}$ subject to the condition*

$$\frac{n_1 + 2}{(m_1 + 1)n_1 + 2m_1 + 1} + \frac{n_2 + 2}{(m_2 + 1)n_2 + 2m_2 + 1} + \frac{1}{q} > 0.$$

Moreover, assume that m_1, n_1, m_2, n_2, q were chosen so that $K = M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ is a knot (see Proposition 4.1). Then K is not smoothly slice.

The conditions on m_i, n_i, q are realized on a multitude of examples. We only list a few but the interested reader will have little trouble finding many additional examples.

Example 1.2. Choose $q = -3$, $m_1 = 1$, and $m_2 = 2$. Then the conditions from Theorem 1.1 translate into

$$n_1, n_2 \geq 1 \quad \text{and} \quad n_1 \text{ is even or } n_2 \text{ is odd.}$$

Consequently there are infinitely many examples among $M(0; [2, n_1 + 2], [3, n_2 + 2], -3)$ which are obstructed from being smoothly slice by Theorem 1.1.

Before stating our next example, we remind the reader that the knot signature $\sigma(K)$ and knot determinant $\det(K)$ of a knot K can be used as obstructions to sliceness. If K is slice then $\sigma(K) = 0$ and $|\det(K)|$ is a square. With this in mind, consider the next example.

Theorem 1.3. *The family $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ (with the restrictions on m_i , n_i and q as outlined in Theorem 1.1) contains an infinite subfamily of knots whose signatures are 0 and whose determinants are square.*

1.1. Organization. In Section 2, we formally outline the approach employed by Lisca to obstruct smooth sliceness. In Section 3, we give definitions of a Montesinos knot as well as outline plumbing 4-manifolds whose boundaries are 2-fold branched covers of such knots. Finally, in Sections 4 and 5, we prove Theorems 1.1 and 1.3 respectively.

2. OBSTRUCTION TO SLICENESS

This section outlines the technique used by Lisca [Lis1]. To start with, let K be a knot in S^3 and let Y_K denote the 3-manifold obtained by taking the double branched cover of S^3 with branching set K . We note that Y_K is always a rational homology sphere and moreover, if K is slice¹, then Y_K bounds a rational homology 4-ball W_K . The manifold W_K is obtained by taking a double branched cover of the 4-ball branched over the slice disk of K .

If K is a knot whose double branched cover Y_K bounds a negative definite smooth 4-manifold X_K , one can form a closed, smooth, negative definite 4-manifold by gluing W_K to X_K along Y_K . Recall Donaldson's celebrated theorem

Theorem 2.1 (Donaldson's Theorem A [Don]). *Let W be a closed, oriented, smooth 4-manifold with negative definite intersection form Q_W , then Q_W is isomorphic to the standard negative definite form $-I$ of the same rank.*

Combining Donaldson's theorem with the remarks preceding it, we arrive at

Theorem 2.2. *If K is a slice knot whose double branched cover Y_K bounds the negative definite smooth 4-manifold X_K , then the intersection form of X_K embeds into the standard negative definite form of the same dimension.*

Example 2.3. Consider K the figure-eight knot. Y_K bounds a plumbing 4-manifold given by a single component of the plumbing given in Figure 1. If K were slice, then there would be an embedding (see Section 4.2 for notation) $\varphi : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ such that $\varphi(f_1)^2 = -3$. No such embedding exists since there are only two basis elements in the codomain. So, Theorem 2.2 implies that K could not be slice.

We know that $K \# K$ is slice. $Y_{K \# K}$ bounds the plumbing 4-manifold in Figure 1. With the indicated basis, we can find an embedding $\varphi : \mathbb{Z}^4 \rightarrow \mathbb{Z}^4$:

$$\varphi(f_1) = e_2 + e_3 + e_4, \quad \varphi(f_2) = e_1 - e_2, \quad \varphi(f_3) = e_1 + e_2 - e_3, \quad \varphi(f_4) = e_3 - e_4.$$

Then, as expected, we cannot use Theorem 2.2 to obstruct $K \# K$ from being slice.



FIGURE 1. Plumbing 4-manifold, whose boundary is the 2-fold cover of $K \# K$, with our chosen basis for its intersection form.

¹All instances of sliceness in the article will always refer to smooth sliceness.

3. MONTESINOS KNOTS AND THEIR 2-FOLD COVERS

Here, we outline the definition of a Montesinos knot and its 2-fold branched cover. We follow [OS]. Let $n_1, n_2, \dots, n_k \in \mathbb{Z}$, then we let $[n_1, n_2, \dots, n_k]$ be the continued fraction

$$[n_1, n_2, \dots, n_k] = n_1 - \frac{1}{n_2 - \frac{1}{\dots - \frac{1}{n_k}}}.$$

Let $\alpha/\beta \in \mathbb{Q}$ be in lowest terms and let $[a_1, a_2, \dots, a_k]$ be a continued fraction representing α/β , then a rational tangle corresponding to α/β is given by Figure 2 where $\boxed{a_i}$ indicates a_i half-twists. With these conventions in place, we can state the definition of a Montesinos link




FIGURE 2. Rational tangles corresponding to $\alpha/\beta = [a_1, a_2, \dots, a_k]$ for k odd and even respectively.

and describe its 2-fold cover.

Definition 3.1. Let $\alpha_1/\beta_1, \alpha_2/\beta_2, \dots, \alpha_n/\beta_n \in \mathbb{Q}$ each be in lowest terms and let $e \in \mathbb{Z}$. Then the Montesinos link $M(e; \alpha_1/\beta_1, \dots, \alpha_n/\beta_n)$ is given by Figure 3 where $\boxed{e}$ represents e half-twists and $\boxed{\alpha_i/\beta_i}$ represent a rational tangle corresponding to α_i/β_i .




FIGURE 3. Montesinos link $M(e; \alpha_1/\beta_1, \dots, \alpha_n/\beta_n)$.

Theorem 3.2 ([Mon]). *The 2-fold branched cover of S^3 branched along a Montesinos link $M(e; \alpha_1/\beta_1, \dots, \alpha_n/\beta_n)$ is the boundary of the plumbing 4-manifold given by Figure 4 where $\alpha_i/\beta_i = [a_1^i, a_2^i, \dots, a_{\ell_i}^i]$.*

Neumann and Raymond [NR] give the following useful result to determine when such a plumbing could be blown down to a negative definite 4-manifold.

Theorem 3.3 ([NR]). *A plumbing, W , with no nonnegative weights in its plumbing graph and whose boundary is the 2-fold branched cover of S^3 branched over the Montesinos link $M(e; \alpha_1/\beta_1, \dots, \alpha_n/\beta_n)$, is negative definite if and only if*

$$e + \sum_{i=1}^n \frac{\beta_i}{\alpha_i} > 0.$$




FIGURE 4. Plumbing 4-manifold X_K with $\partial X_K = Y_K$ for $K = M(e; \alpha_1/\beta_1, \dots, \alpha_n/\beta_n)$.

4. OBSTRUCTING SLICENESS IN $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$

In this section, we consider $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ and prove Theorem 1.1. First, we outline the restrictions on m_i, n_i, q found within Theorem 1.1. In order for $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ to be a knot we need to carefully restrict the choices of parity on m_i, n_i, q . The following proposition outlines these restrictions.

Proposition 4.1. *All but the following twelve combinations of parity for the m_i , n_i , and q , result in knots in the family $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$:*

- (1) q , m_1 , and $n_1 + 1$ even and no other restrictions,
- (2) q , m_2 , and $n_2 + 1$ even and no other restrictions,
- (3) q , m_1 , n_1 , $n_2 + 1$ odd and no other restrictions,
- (4) q , $n_1 + 1$, m_2 , n_2 odd and no other restrictions,
- (5) q , $m_1 + 1$, n_1 , $m_2 + 1$, and n_2 odd.

Proof. The proof is a simple matter of an exhaustive check of the thirty-two possible combinations of parity for the m_i , n_i and q . $\square$

To apply Theorem 2.2, we need the 2-fold cover of $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ to bound a negative definite 4-manifold. In light of Theorem 3.3, we require that m_i , n_i and q satisfy

$$\frac{n_1 + 2}{(m_1 + 1)n_1 + 2m_1 + 1} + \frac{n_2 + 2}{(m_2 + 1)n_2 + 2m_2 + 1} + \frac{1}{q} > 0. \quad (1)$$

4.1. Choosing a Plumbing 4-manifold. Rather than applying Theorem 3.2 directly, we note that $[m + 1, n + 2]$ can be expressed as a different continued fraction expansion - one which is more compatible with Lisca's approach.

Proposition 4.2. *Let $m, n \in \mathbb{N}$, then $[m + 1, n + 2] = [-1, -1, \underbrace{-2, \dots, -2}_m, -3, \underbrace{-2, \dots, -2}_n]$.*

Proof. This fact is easily proven by induction. $\square$

This, combined with Theorem 3.2, gives that the 2-fold cover of $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ bounds the plumbing 4-manifold in Figure 5. Blowing down appropriate vertices results in the negative definite 4-manifold in Figure 6.

4.2. Proving Theorem 1.1. Before we present the proof of Theorem 1.1, we define our notation for embeddings of intersection forms over the integers. For an integer n , we view the $n \times n$ matrix A as an intersection pairing on $\mathbb{Z}^n$ with respect to the standard basis $f_1, \dots, f_n$. Similarly, we view $-I$ as the standard negative definite intersection pairing on another copy




FIGURE 5. Plumbing 4-manifold whose boundary is Y_K for $K = M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$.

of $\mathbb{Z}^n$ with respect to its standard basis, $e_1, \dots, e_n$. We abbreviate $\langle Af_i, f_j \rangle$ to $f_i \cdot f_j$, and similarly for $-I$. Finally, an embedding of A into $-I$ is a monomorphism $\varphi : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ such that $f_i \cdot f_j = \varphi(f_i) \cdot \varphi(f_j)$.

Proof of Theorem 1.1. Let Q_W be the intersection form associated to the plumbing 4-manifold in Figure 6, we show that Q_W does not embed into the standard negative definite form of equal rank. Suppose to the contrary that Q_W does embed into $-I$. Then, there must exist an




FIGURE 6. Plumbing 4-manifold with our chosen basis for Q_W .

embedding $\varphi : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$, where $n = n_1 + m_1 + m_2 + n_2 + 2$. Taking the f_i indicated in Figure 6 as our basis for $Q_W = [q_{ij}]$, we explore the structure of φ .

First, note that up to a change of basis for $-I$, any vertex with self intersection -2 or -3 has essentially a unique image under φ . For instance, we can assume that, $\varphi(f_1) = e_1 - e_2$. Similarly, we can assume that $\varphi(f_2) = e_i - e_{i+1}$ for some $i \in \{1, 2, \dots, n\}$ (up to a change of basis). Now $f_1 \cdot f_2 = q_{12} = 1$; therefore $\varphi(f_1) \cdot \varphi(f_2) = 1$. By definition

$$\varphi(f_1) \cdot \varphi(f_2) = e_1 \cdot e_i - e_1 \cdot e_{i+1} - e_2 \cdot e_i + e_2 \cdot e_{i+1}.$$

Since φ is injective, we cannot have $i = 1$. If $i > 2$, then the above quantity is zero, so it follows that $i = 2$ and thus $\varphi(f_2) = e_2 - e_3$. Similar arguments show that

$$\varphi(f_i) = e_i - e_{i+1}, \quad 1 \leq i \leq n_1. \quad (2)$$

It is worth noting that if $n_1 > 2$, we could take $\varphi(f_3) = -e_1 - e_2$ while still satisfying the pairings $f_1 \cdot f_3 = 0$ and $f_2 \cdot f_3 = 1$. However, with such a definition, we still need $f_4 \cdot f_3 = 1$, implying that $\varphi(f_4)$ must contain either $\pm e_1$ or $\pm e_2$, but not both. However, this would imply that $\varphi(f_1)$ would pair with $\varphi(f_4)$ nontrivially - which contradicts the fact that φ is an embedding. It follows that, up to a change of basis for $-I$, the images defined in (2) are the only possibilities.

Next, we consider the image of f_{n_1+2} under φ . As above, up to a change of basis we can take $\varphi(f_{n_1+2}) = e_{n_1+2} - e_{n_1+3}$ and by the same argument used for f_2 , we arrive at

$$\varphi(f_i) = e_i - e_{i+1}, \quad n_1 + 2 \leq i \leq n_1 + m_1 + m_2.$$

Similarly, we can assume that $\varphi(f_{n_1+m_1+m_2+2}) = e_{n_1+m_1+m_2+2} - e_{n_1+m_1+m_2+3}$ and thus

$$\varphi(f_i) = e_i - e_{i+1}, \quad n_1 + m_1 + m_2 + 2 \leq i \leq n_1 + m_1 + m_2 + n_2 + 1.$$

Next, we consider $\varphi(f_{n_1+1})$. Since φ is a homomorphism, we know that $\varphi(f_{n_1+1}) = \sum_{i=1}^{n_1+m_1+m_2+n_2+2} \lambda_i e_i$ for some $\lambda_i \in \mathbb{Z}$. Given that $f_i \cdot f_{n_1+1} = \varphi(f_i) \cdot \varphi(f_{n_1+1})$, we have

$$\varphi(f_i) \cdot \varphi(f_{n_1+1}) = \begin{cases} \pm 1 & i = n_1, \\ -3 & i = n_1 + 1, \\ 1 & i = n_1 + 2, \\ 0 & \text{otherwise.} \end{cases} \quad (3)$$

Note that the ± 1 (rather than just 1) arises from the fact that we may have had to change the basis of $-I$ to get images of the previous vertices in their correct forms. Thus, we may have caused the pairing of $\varphi(f_{n_1})$ and $\varphi(f_{n_1+1})$ to become negative.

Since $\varphi(f_i) = e_i - e_{i+1}$ for each $i \neq n_1 + 1, n_1 + m_1 + m_2 + 1, n_1 + m_1 + m_2 + n_2 + 2$, we have that $\varphi(f_i) \cdot \varphi(f_{n_1+1}) = \lambda_{i+1} - \lambda_i$ for the same i . Then, from (3) we have that

$$\lambda_{i+1} - \lambda_i = 0,$$

for $1 \leq i \leq n_1 - 1$, $n_1 + 3 \leq i \leq n_1 + m_1 + m_2$, and $n_1 + m_1 + m_2 + 3 \leq n_1 + m_1 + m_2 + n_2 + 1$. It follows that

$$\lambda_1 = \lambda_2 = \dots = \lambda_{n_1}, \quad \lambda_{n_1+3} = \lambda_{n_1+4} = \dots = \lambda_{n_1+m_1+m_2+1},$$

$$\lambda_{n_1+m_1+m_2+3} = \lambda_{n_1+m_1+m_2+4} = \dots = \lambda_{n_1+m_1+m_2+n_2+2}.$$

Let $\lambda = \lambda_1$, $\eta = \lambda_{n_1+3}$ and $\gamma = \lambda_{n_1+m_1+m_2+3}$. Now, $\varphi(f_{n_1}) \cdot \varphi(f_{n_1+1}) = \lambda_{n_1} - \lambda_{n_1+1} = \lambda_{n_1} - \lambda = \pm 1$. Therefore, $\lambda_{n_1} = \lambda \pm 1$. Similarly, $\lambda_{n_1+2} = \eta - 1$. So,

$$\begin{aligned} \varphi(f_{n_1+1}) &= \\ &= \sum_{i=1}^{n_1} \lambda e_i + (\lambda \pm 1) e_{n_1+1} + (\eta - 1) e_{n_1+2} + \sum_{i=n_1+3}^{n_1+m_1+m_2+1} \eta e_i + \sum_{i=n_1+m_1+m_2+2}^{n_1+m_1+m_2+n_2+2} \gamma e_i. \end{aligned}$$

Direct calculation along with (3) gives that

$$\varphi(f_{n_1+1})^2 = - \left(n_1 \lambda^2 + (\lambda \pm 1)^2 + (\eta - 1)^2 + (m_1 + m_2 - 1) \eta^2 + (n_2 + 1) \gamma^2 \right) = -3.$$

Clearly, if $n_2 > 2$, then $\gamma = 0$. If $n_2 = 2$ and $\gamma \neq 0$, then λ^2 , $(\lambda \pm 1)^2$, $(\eta - 1)^2$, and η^2 must be identically zero - clearly an impossibility; therefore if $n_2 = 2$, $\gamma = 0$. Moreover, if $n_2 = 1$ and $\gamma \neq 0$, then three of λ^2 , $(\lambda \pm 1)^2$, $(\eta - 1)^2$, and η^2 must be zero - which is again impossible. It follows that γ is necessarily 0. Therefore

$$\varphi(f_{n_1+1}) = \sum_{i=1}^{n_1} \lambda e_i + (\lambda \pm 1) e_{n_1+1} + (\eta - 1) e_{n_1+2} + \sum_{i=n_1+3}^{n_1+m_1+m_2+1} \eta e_i.$$

The same argument applies to $\varphi(f_{n_1+m_2+m_2+1})$ where the pairings are given by

$$\varphi(f_i) \cdot \varphi(f_{n_1+m_1+m_2+1}) = \begin{cases} \pm 1 & i = n_1 + m_1 + m_2 + 2, \\ -3 & i = n_1 + m_1 + m_2 + 1, \\ 1 & i = n_1 + m_1 + m_2, \\ 0 & \text{otherwise.} \end{cases} \quad (4)$$

Then, there exist $\alpha, \beta \in \mathbb{Z}$ such that

$$\begin{aligned} \varphi(f_{n_1+m_1+m_2+1}) &= \\ &= \sum_{i=n_1+2}^{n_1+m_1+m_2} \beta e_i + (\beta+1) e_{n_1+m_1+m_2+1} + (\alpha \pm 1) e_{n_1+m_1+m_2+2} + \sum_{i=n_1+m_1+m_2+3}^{n_1+m_1+m_2+n_2+2} \alpha e_i. \end{aligned}$$

Therefore, we have explicit forms for the images of all but the last basis element in Q_W under φ . Before we consider this last image, we extract as much information out of the pairings in (3) and (4) as we can. (4) gives that $\varphi(f_{n_1+1}) \cdot \varphi(f_{n_1+m_1+m_2+1}) = 0$. Therefore,

$$\varphi(f_{n_1+1}) \cdot \varphi(f_{n_1+m_1+m_2+1}) = -(\beta(\eta-1) + (m_1+m_2-2)\beta\eta + (\beta+1)\eta) = 0$$

and thus

$$(m_1+m_2)\beta\eta - \beta + \eta = 0.$$

Noting that the coefficient of each basis element in the image of a square -3 vertex is necessarily in the set $\{0, -1, 1\}$, we have that $\beta, \beta+1, \eta, \eta-1 \in \{0, -1, 1\}$; so, $\beta = 0$ or $\beta = -1$ and $\eta = 0$ or $\eta = 1$.

If $\beta = 0$, then $\eta = 0$ and if $\beta = -1$, then $\eta = 1$ and $m_1+m_2 = 2$. In this latter case, α is forced to be ± 1 which, in turn, forces $n_2 = 2$. Moreover, λ is forced to be ± 1 which makes $n_1 = 2$. We'll come back to this case.

Now, suppose that $\beta = \eta = 0$, then

$$\varphi(f_{n_1+1}) = \sum_{i=1}^{n_1} \lambda e_i + (\lambda \pm 1) e_{n_1+1} - e_{n_1+2},$$

$$\varphi(f_{n_1+m_1+m_2+1}) = e_{n_1+m_1+m_2+1} + (\alpha \pm 1) e_{n_1+m_1+m_2+2} + \sum_{i=n_1+m_1+m_2+3}^{n_1+m_1+m_2+n_2+2} \alpha e_i.$$

Moreover, $\lambda = \mp 1$ and $n_1 = 2$, $\alpha = \mp 1$ and $n_2 = 2$, so

$$\varphi(f_{n_1+1}) = \varphi(f_3) = \mp e_1 \mp e_2 - e_4,$$

$$\varphi(f_{n_1+m_1+m_2+1}) = \varphi(f_{m_1+m_2+3}) = e_{m_1+m_2+3} \mp e_{m_1+m_2+5} \mp e_{m_1+m_2+6}.$$

Therefore, if n_1 and n_2 are not identically 2, then no such embedding φ exists.

Corresponding to the case when $\beta = \eta = 0$, we have $n_1 = n_2 = 2$ and $m_1, m_2 \in \mathbb{N}$. Then, we know that up to a change of basis,

$$\begin{aligned} \varphi(f_1) &= e_1 - e_2 \\ \varphi(f_2) &= e_2 - e_3 \\ \varphi(f_3) &= \mp e_1 \mp e_2 - e_4 \\ \varphi(f_4) &= e_4 - e_5 \\ &\vdots \\ \varphi(f_{m_1+m_2+2}) &= e_{m_1+m_2+3} - e_{m_1+m_2+2} \\ \varphi(f_{m_1+m_2+3}) &= e_{m_1+m_2+3} \mp e_{m_1+m_2+5} \mp e_{m_1+m_2+6} \\ \varphi(f_{m_1+m_2+4}) &= e_{m_1+m_2+5} - e_{m_1+m_2+4} \\ \varphi(f_{m_1+m_2+5}) &= e_{m_1+m_2+6} - e_{m_1+m_2+5} \end{aligned}$$

Now, suppose that $\varphi(f_{m_1+m_2+6}) = \sum_{i=1}^{m_1+m_2+6} \mu_i e_i$ for $\mu_i \in \mathbb{Z}$. Then, we get the following system of equations arising from the pairings indicated in Figure 6

$$\left\{ \begin{array}{l} \varphi(f_1) \cdot \varphi(f_{m_1+m_2+6}) = \mu_2 - \mu_1 = 0 \\ \varphi(f_2) \cdot \varphi(f_{m_1+m_2+6}) = \mu_3 - \mu_2 = 0 \\ \varphi(f_i) \cdot \varphi(f_{m_1+m_2+6}) = \mu_{i+1} - \mu_i = 0 \quad 4 \leq i \leq m_1 + 2, \quad m_1 + 4 \leq i \leq m_1 + m_2 + 2 \\ \varphi(f_{m_1+m_2+4}) \cdot \varphi(f_{m_1+m_2+6}) = \mu_{m_1+m_2+5} - \mu_{m_1+m_2+4} = 0 \\ \varphi(f_{m_1+m_2+5}) \cdot \varphi(f_{m_1+m_2+6}) = \mu_{m_1+m_2+6} - \mu_{m_1+m_2+5} = 0 \\ \varphi(f_{m_1+3}) \cdot \varphi(f_{m_1+m_2+6}) = \mu_{m_1+4} - \mu_{m_1+3} = 1 \\ \varphi(f_3) \cdot \varphi(f_{m_1+m_2+6}) = \pm\mu_1 \pm \mu_2 + \mu_4 = 0 \\ \varphi(f_{m_1+m_2+3}) \cdot \varphi(f_{m_1+m_2+6}) = -\mu_{m_1+m_2+3} \pm \mu_{m_1+m_2+5} \pm \mu_{m_1+m_2+6} = 0 \end{array} \right.$$

Then $\mu_1 = \mu_2 = \mu_3$, $\mu_4 = \mu_5 = \dots = \mu_{m_1+3}$, $\mu_{m_1+4} = \mu_{m_1+5} = \dots = \mu_{m_1+m_2+3}$, and $\mu_{m_1+m_2+4} = \mu_{m_1+m_2+5} = \mu_{m_1+m_2+6}$. Let $\mu = \mu_1$, $\nu = \mu_4$, $\rho = \mu_{m_1+4}$ and $\sigma = \mu_{m_1+m_2+4}$, then

$$\begin{aligned} \varphi(f_{m_1+m_2+6}) &= \\ &= \mu(e_1 + e_2 + e_3) + \sum_{i=4}^{m_1+3} \nu e_i + \sum_{i=m_1+4}^{m_1+m_2+3} \rho e_i + \sigma(e_{m_1+m_2+4} + \dots + e_{m_1+m_2+6}) \end{aligned}$$

By construction, the above satisfies the first five equations in the system. Since $e_{m_1+3} - e_{m_1+4} = 1$, we have that $\nu = \rho + 1$; now, $\pm\mu_1 \pm \mu_2 + \mu_4 = 0$ gives that $\mp 2\mu = \nu$ and thus ν is necessarily even which means that ρ is odd; however, $-\mu_{m_1+m_2+3} \pm \mu_{m_1+m_2+5} \pm \mu_{m_1+m_2+6} = 0$ gives that $\rho = \pm 2\sigma$ - a contradiction. Therefore no such function φ could exist if $\beta = \eta$.

It follows that the only way for such φ to have any hope of existing is if we fall into the latter case mentioned above. In this case, $m_1 = m_2 = 1$, $n_1 = n_2 = 2$, $\beta = -1$, $\eta = 1$, $\alpha = \pm 1$, and $\lambda = \pm 1$. Then, again, up to a change of basis, we can take

$$\begin{aligned} \varphi(f_1) &= e_1 - e_2 \\ \varphi(f_2) &= e_2 - e_3 \\ \varphi(f_3) &= \pm e_1 \pm e_2 + e_5 \\ \varphi(f_4) &= e_4 - e_5 \\ \varphi(f_5) &= -e_4 \pm e_7 \pm e_8 \\ \varphi(f_6) &= e_6 - e_7 \\ \varphi(f_7) &= e_7 - e_8 \end{aligned}$$

Suppose that $\varphi(f_8) = \sum_{i=1}^8 \mu_i e_i$ for $\mu_i \in \mathbb{Z}$. As above, we get the following system of equations arising from the pairings indicated in Figure 6


$$\begin{aligned} \varphi(f_1) \cdot \varphi(f_8) &= \mu_2 - \mu_1 = 0 & \varphi(f_5) \cdot \varphi(f_8) &= \mu_4 \mp \mu_7 \mp \mu_8 = 0 \\ \varphi(f_2) \cdot \varphi(f_8) &= \mu_3 - \mu_2 = 0 & \varphi(f_6) \cdot \varphi(f_8) &= \mu_7 - \mu_6 = 0 \\ \varphi(f_3) \cdot \varphi(f_8) &= \mp \mu_1 \mp \mu_2 - \mu_5 = 0 & \varphi(f_7) \cdot \varphi(f_8) &= \mu_8 - \mu_7 = 0 \\ \varphi(f_4) \cdot \varphi(f_8) &= \mu_5 - \mu_4 = 1 \end{aligned}$$

Let $\mu = \mu_1$ and $\nu = \mu_6$, then $\mu = \mu_1 = \mu_2 = \mu_3$ and $\nu = \mu_6 = \mu_7 = \mu_8$. So $\mu_4 = \pm 2\nu$ and $\mu_5 = \mp 2\mu$. But, $\mu_5 = 1 + \mu_4$, which means that $\mp 2\mu = 1 \pm 2\nu$ - a contradiction. So no such function φ exists in this case either.

Given that we have exhausted all possible cases, it is clear that no embedding $\varphi : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ could exist for Q_W arising from the plumbing indicated in Figure 6. It follows that Q_W does not embed into the standard negative definite form $-I$ as claimed. Then, the result follows from Theorem 2.2. Therefore, no member of $M([m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ with the outlined restrictions on m_i, n_i, q can be smoothly slice. $\square$

5. THE INFINITE SUBFAMILY

Proof of Theorem 1.3. Consider the case with $m_1 = n_2 = 1$, $m_2 = 2$, and $q = -3$. Then, we show that $M(0; [2, n_1 + 2], [3, 3], -3)$ contains our desired subfamily. With these choices of m_i , n_2 , and q , we can take any choice of n_1 to meet both the requirement for negative definiteness and the requirements on parity. Thus every knot $M(0; [2, n_1 + 2], [3, 3], -3)$ meets the restrictions outlined in Theorem 1.1.


 FIGURE 7. $M(0; [2, n_1 + 2], [3, 3], -3)$

Now, we use the method outlined in [GL] to calculate the signature and determinant of each member of $M(0; [2, n_1 + 2], [3, 3], -3)$. We begin by checkerboard coloring the knot diagram in Figure 7 with the unbounded exterior region colored white. From this coloring, we arrive at the symmetric Goeritz matrix given by

$$G = \begin{bmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 0 & 0 & -3 \\ 0 & 1 & 0 & 0 & 0 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & n_1 + 1 & -n_1 - 2 \\ 0 & 0 & 0 & 0 & -3 & 0 & -n_1 - 2 & n_1 + 6 \end{bmatrix}$$

Direct calculation gives that $\det(G) = 26n_1 + 51$, which as [GL] shows, is precisely the Determinant of the knot $M(0; [2, n_1 + 2], [3, 3], -3)$. Summing over all type II crossings in Figure 7 with our chosen coloring scheme and an arbitrary string orientation gives a correction factor of $\mu(M(0; [2, n_1 + 2], [3, 3], -3)) = 3 - 3 = 0$. Therefore, the signature of $M(0; [2, n_1 + 2], [3, 3], -3)$ is precisely the signature of G . It's easy to verify that the signature of G is 0 for any valid choice of n_1 and thus each member of $M(0; [2, n_1 + 2], [3, 3], -3)$ has signature 0.

Therefore, each n_1 such that $26n_1 + 51 = a^2$ for some $a \in \mathbb{N}$, gives an example of a knot in the family $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ which has square determinant and signature 0. To show that this is an infinite subset, consider the sequence $\{a_n\}_{n=0}^{\infty}$ given recursively by

$a_0 = 21$, and

$$a_{n+1} = \begin{cases} a_n + 10 & \text{if } n \text{ is even,} \\ a_n + 16 & \text{if } n \text{ is odd.} \end{cases}$$

We claim that there is a positive integral solution, n_1 , to $26n_1 + 51 = a_n^2$ for each n . To see this, note that $26 \cdot 15 + 51 = 441 = a_0^2$ and $26 \cdot 35 + 51 = 961 = a_1^2$ and suppose that for $k = n - 1, n$, there exists an $\eta \in \mathbb{N}$ such that $26\eta + 51 = a_k^2$. Then, consider $26\eta + 51 = a_{n+1}^2$. By definition $a_{n+1} = a_{n-1} + 26$, therefore

$$26\eta + 51 = a_{n-1}^2 + 52a_{n-1} + 26^2$$

Solving for a_{n-1}^2 gives

$$26(\eta - 2a_{n-1} - 26) + 51 = a_{n-1}^2.$$

By induction, this has a solution in $\mathbb{N}$, call it ℓ . Then, we can take $\eta = \ell + 2a_{n-1} + 26$. It follows that each a_n in $\{a_n\}_{n=0}^\infty$ admits a distinct positive integer valued solution to $26\eta + 51 = a_n^2$ and thus $M(0; [m_1 + 1, n_1 + 2], [m_2 + 1, n_2 + 2], q)$ (with the restrictions outlined in Theorem 1.1) contains an infinite subfamily of knots whose signatures are 0 and whose determinants are square. $\square$

REFERENCES

- [Don] S. K. Donaldson. The orientation of Yang-Mills moduli spaces and 4-manifold topology. *J. Differential Geom.*, 26(3):397–428, 1987.
- [GJ] Joshua Greene and Stanislav Jabuka. The slice-ribbon conjecture for 3-stranded pretzel knots, 2007.
- [GL] C. McA. Gordon and R. A. Litherland. On the signature of a link. *Invent. Math.*, 47(1):53–69, 1978.
- [Kir] Rob Kirby. Problems in low dimensional manifold theory. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2*, Proc. Sympos. Pure Math., XXXII, pages 273–312. Amer. Math. Soc., Providence, R.I., 1978.
- [Lis1] Paolo Lisca. Lens spaces, rational balls and the ribbon conjecture. *Geom. Topol.*, 11:429–472, 2007.
- [Lis2] Paolo Lisca. Sums of lens spaces bounding rational balls. *Algebr. Geom. Topol.*, 7:2141–2164, 2007.
- [Mon] José M. Montesinos. Seifert manifolds that are ramified two-sheeted cyclic coverings. *Bol. Soc. Mat. Mexicana (2)*, 18:1–32, 1973.
- [NR] Walter D. Neumann and Frank Raymond. Seifert manifolds, plumbing, μ -invariant and orientation reversing maps. In *Algebraic and geometric topology (Proc. Sympos., Univ. California, Santa Barbara, Calif., 1977)*, volume 664 of *Lecture Notes in Math.*, pages 163–196. Springer, Berlin, 1978.
- [OS] Brendan Owens and Sašo Strle. Rational homology spheres and the four-ball genus of knots. *Adv. Math.*, 200(1):196–216, 2006.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824
E-mail address: will2086@msu.edu